
Leader in Biomolecular Solutions for Life Science



Catalog No.: A21085 4 Publications



## **Basic Information**

Observed MW 48kDa

Calculated MW 46kDa

Category Primary antibody

Applications WB,ELISA

Cross-Reactivity Mouse

CloneNo number ATC0001

## Background

Enables cysteine-type endopeptidase activity. Involved in several processes, including positive regulation of I-kappaB kinase/NF-kappaB signaling; positive regulation of interleukin-1 beta production; and protein autoprocessing. Acts upstream of or within several processes, including membrane hyperpolarization; mitochondrial depolarization; and positive regulation of interleukin-1 alpha production. Located in extracellular region. Part of IPAF inflammasome complex. Is expressed in knee joint and metanephros. Orthologous to human CASP1 (caspase 1).

# Immunogen Information

# WB 1:1000 - 1:3000 ELISA Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

**Recommended Dilutions** 

Gene ID 12362 Swiss Prot P29452

### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms ICE; II1bc

## Contact

6	400-999-6126
$\mathbf{X}$	cn.market@abclonal.com.cn
€	www.abclonal.com.cn

## **Product Information**

**Source** Rat **lsotype** lgG **Purification** Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

