

Acetyl-Histone H3-K9 Rabbit mAb

Catalog No.: A21107 **Recombinant** **1 Publications**

Basic Information

Observed MW

17kDa

Calculated MW

15kDa

Category

Primary antibody

Applications

ELISA,DB,WB,IHC-
P,IF/ICC,IP,ChIP,CUT&Tag,ChIP-seq

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range
Predicted)

CloneNo number

ARC50576

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

DB 1:500 - 1:1000**WB** 1:500 - 1:1000**IHC-P** 1:100 - 1:500**IF/ICC** 1:50 - 1:200**IP** 0.5µg-4µg antibody for
200µg-400µg extracts of
whole cells**ChIP** 5µg antibody for
5µg-10µg of Chromatin**CUT&Tag** 10⁵ cells /1 µg**ChIP-seq** 1:50 - 1:100

Immunogen Information

Gene ID

8290/8350

Swiss Prot

Q16695/P68431

Immunogen

A synthetic acetylated peptide around K9 of human Histone H3 (NP_003520.1).

Synonyms

H3/A; H3C2; H3C3; H3C4; H3C6; H3C7; H3C8; H3FA; H3C10; H3C11; H3C12; HIST1H3A;
Acetyl-Histone H3-K9

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data




Western blot analysis of various lysates using Acetyl-Histone H3-K9 Rabbit mAb (A21107) at 1:1000 dilution. HeLa and NIH/3T3 and C6 cells were treated by TSA (1 μ M) at 37°C for 18 hours. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.



Immunoprecipitation analysis of 600 μ g extracts of NIH/3T3 cells using 5 μ g Acetyl-Histone H3-K9 antibody (A21107). Western blot was performed from the immunoprecipitate using Acetyl-Histone H3-K9 antibody (A21107) at a dilution of 1:1000.




Chromatin immunoprecipitation analysis of extracts of HeLa cells, using Acetyl-Histone H3-K9 Rabbit mAb (A21107) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.






Chromatin immunoprecipitations were performed with cross-linked chromatin from 293F cells and Acetyl-Histone H3-K9 (A21107). The ChIP sequencing results indicate the enrichment pattern of Acetyl-Histone H3-K9 in selected genomic region and representative gene loci (RPL30), as shown in figure.

Validation Data




Chromatin immunoprecipitations were performed with cross-linked chromatin from 293F cells and Acetyl-Histone H3-K9 (A21107). The ChIP sequencing results indicate the enrichment pattern of Acetyl-Histone H3-K9 in selected genomic region and representative gene loci (RPL30), as shown in figure.




CUT&Tag was performed using the CUT&Tag Assay Kit (pAG-Tn5) for Illumina(RK20265) from 10⁵ K562 cells with 1 µg Acetyl-Histone H3-K9 Rabbit mAb antibody (A21107), along with a Goat Anti-Rabbit IgG(H+L). The CUT&Tag results indicate the enrichment pattern of Acetyl-Histone H3-K9 in representative gene loci (RPL30), as shown in figure.

Dot-blot analysis of all sorts of peptides using Acetyl-Histone H3-K9 antibody (A21107) at 1:1000 dilution.


Confocal imaging of HeLa TSA and HeLa cells using Acetyl-Histone H3-K9 Rabbit mAb (A21107,dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007,dilution 1:500)(Red).DAPI was used for nuclear staining (Blue). Objective: 100x.




Immunohistochemistry analysis of Acetyl-Histone H3-K9 in paraffin-embedded human colon carcinoma using Acetyl-Histone H3-K9 Rabbit mAb (A21107) at dilution of 1:200 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of Acetyl-Histone H3-K9 in paraffin-embedded human liver using Acetyl-Histone H3-K9 Rabbit mAb (A21107) at dilution of 1:200 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of Acetyl-Histone H3-K9 in paraffin-embedded human spleen using Acetyl-Histone H3-K9 Rabbit mAb (A21107) at dilution of 1:200 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of Acetyl-Histone H3-K9 in paraffin-embedded mouse brain using Acetyl-Histone H3-K9 Rabbit mAb (A21107) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of Acetyl-Histone H3-K9 in paraffin-embedded mouse intestine using Acetyl-Histone H3-K9 Rabbit mAb (A21107) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of Acetyl-Histone H3-K9 in paraffin-embedded mouse spleen using Acetyl-Histone H3-K9 Rabbit mAb (A21107) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of Acetyl-Histone H3-K9 in paraffin-embedded mouse testis using Acetyl-Histone H3-K9 Rabbit mAb (A21107) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of Acetyl-Histone H3-K9 in paraffin-embedded rat brain using Acetyl-Histone H3-K9 Rabbit mAb (A21107) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.