

[KO Validated] YAP1 Rabbit mAb

Catalog No.: A21216 **KO Validated** **Recombinant** **10 Publications**

Basic Information

Observed MW

70kDa

Calculated MW

54kDa

Category

Primary antibody

Applications

WB, IHC-P, IF/ICC, IP, ELISA, ChIP

Cross-Reactivity

Human, Mouse, Rat

Clone/No. number

ARC53479

Background

This gene encodes a downstream nuclear effector of the Hippo signaling pathway which is involved in development, growth, repair, and homeostasis. This gene is known to play a role in the development and progression of multiple cancers as a transcriptional regulator of this signaling pathway and may function as a potential target for cancer treatment. Alternative splicing results in multiple transcript variants encoding different isoforms.

Recommended Dilutions

WB 1:20000 - 1:80000

IHC-P 1:200 - 1:800

IF/ICC 1:200 - 1:800

IP 0.5µg-4µg antibody for 200µg-400µg extracts of whole cells

ELISA Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

ChIP 5µg antibody for 10µg-15µg of Chromatin

Immunogen Information

Gene ID

10413

Swiss Prot

P46937

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

YAP; YKI; COB1; YAP2; YAP-1; YAP65; P1

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Contact

-  | 400-999-6126
-  | cn.market@abclonal.com.cn
-  | www.abclonal.com.cn


Validation Data




Immunoprecipitation analysis of 300 µg extracts of HeLa cells using 3 µg YAP1 antibody (A21216). Western blot was performed from the immunoprecipitate using [KO Validated] YAP1 Rabbit mAb (A21216) at a dilution of 1:2000.




Validation Data




Chromatin immunoprecipitation analysis of extracts of HeLa cells, using [KO Validated] YAP1 Rabbit mAb (A21216) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.




Confocal imaging of C2C12 cells using [KO Validated] YAP1 Rabbit mAb (A21216, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Confocal imaging of HeLa cells using [KO Validated] YAP1 Rabbit mAb (A21216, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Immunohistochemistry analysis of paraffin-embedded human breast cancer tissue using [KO Validated] YAP1 Rabbit mAb (A21216) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded human thyroid cancer tissue using [KO Validated] YAP1 Rabbit mAb (A21216) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded human tonsil tissue using [KO Validated] YAP1 Rabbit mAb (A21216) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded mouse spleen tissue using [KO Validated] YAP1 Rabbit mAb (A21216) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.

Validation Data



Immunohistochemistry analysis of paraffin-embedded rat spleen tissue using [KO Validated] YAP1 Rabbit mAb (A21216) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.