

MonoMethyl-Histone H3-K79 Rabbit mAb

Catalog No.: A21223 **Recombinant**

Basic Information

Observed MW

17kDa

Calculated MW

16kDa

Category

Primary antibody

Applications

WB,DB,IHC-P,ELISA,ChIP,ChIP-seq

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

CloneNo number

ARC53324

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

WB 1:4000 - 1:16000**DB** 1:2000 - 1:20000**IHC-P** 1:1000 - 1:5000**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.**ChIP** 5µg antibody for 5µg-10µg of Chromatin**ChIP-seq** 1:50 - 1:200

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn

Immunogen Information

Gene ID

8290/8350

Swiss Prot

Q16695/P68431

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

H3/A; H3C2; H3C3; H3C4; H3C6; H3C7; H3C8; H3FA; H3C10; H3C11; H3C12; HIST1H3A; MonoMethyl-Histone H3-K79

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




www.abclonal.com.cn


Validation Data




Western blot analysis of various lysates, using MonoMethyl-Histone H3-K79 Rabbit mAb (A21223) at 1:4000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.




Chromatin immunoprecipitations were performed with cross-linked chromatin from HeLa cells and MonoMethyl-Histone H3-K79 Rabbit mAb (A21223). The ChIP sequencing results indicate the enrichment pattern of MonoMethyl-Histone H3-K79 in selected genomic region and representative gene loci (RPL30), as shown in figure.




Chromatin immunoprecipitation analysis of extracts of HeLa cells, using MonoMethyl-Histone H3-K79 Rabbit mAb antibody (A21223) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Dot-blot analysis of all sorts of peptides




Immunohistochemistry analysis of paraffin-



Immunohistochemistry analysis of paraffin-


Validation Data

using MonoMethyl-Histone H3-K79 antibody (A21223) at 1:20000 dilution.




Immunohistochemistry analysis of paraffin-embedded Human colon tissue using MonoMethyl-Histone H3-K79 Rabbit mAb (A21223) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.

embedded Human breast tissue using MonoMethyl-Histone H3-K79 Rabbit mAb (A21223) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human breast tissue using MonoMethyl-Histone H3-K79 Rabbit mAb (A21223) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.


embedded Human colon carcinoma tissue using MonoMethyl-Histone H3-K79 Rabbit mAb (A21223) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using MonoMethyl-Histone H3-K79 Rabbit mAb (A21223) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse testis tissue using MonoMethyl-Histone H3-K79 Rabbit mAb (A21223) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using MonoMethyl-Histone H3-K79 Rabbit mAb (A21223) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat testis tissue using MonoMethyl-Histone H3-K79 Rabbit mAb (A21223) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.