Leader in Biomolecular Solutions for Life Science

CSNK2A2 Rabbit pAb

Catalog No.: A21368 1 Publications



Basic Information

Observed MW 41kDa

Calculated MW 41kDa

Category Primary antibody

Applications ELISA,WB

Cross-Reactivity Human, Mouse

Background

This gene encodes the alpha', or alpha 2, catalytic subunit of the protein kinase enzyme, casein kinase 2 (CK2). Casein kinase 2 is a serine/threonine protein kinase that phosphorylates acidic proteins such as casein. It is involved in various cellular processes, including cell cycle control, apoptosis, and circadian rhythms. This heterotetrameric kinase includes two catalytic subunits, either alpha or alpha', and two regulatory beta subunits. The closely related gene paralog encoding the alpha, or alpha 1 subunit (CSNK2A1, Gene ID: 1457) is found on chromosome 20. An intronic variant in this gene (alpha 2) may be associated with leukocyte telomere length in a South Asian population. A related transcribed pseudogene is found on chromosome 11.

Recommended Dilutions

1:500 - 1:2000

Immunogen Information

WB

Gene ID 1459

Swiss Prot P19784

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-350 of human CSNK2A2 (NP_001887.1).

Synonyms

CK2A2; CSNK2A1; CK2alpha'; CSNK2A2


a 400-999-6126 x cn.market@abclonal.com.cn y www.abclonal.com.cn

Product Information

Source Rabbit **Isotype** IgG Purification Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of various lysates, using CSNK2A2 Rabbit pAb (A21368) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 20s.