# ABclonal www.abclonal.com

# Integrin alpha 5 (ITGA5/CD49e) Rabbit pAb

Catalog No.: A22018

# **Basic Information**

#### **Observed MW**

150kDa

#### **Calculated MW**

115kDa

#### Category

Primary antibody

#### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Mouse

# **Background**

The product of this gene belongs to the integrin alpha chain family. Integrins are heterodimeric integral membrane proteins composed of an alpha subunit and a beta subunit that function in cell surface adhesion and signaling. The encoded preproprotein is proteolytically processed to generate light and heavy chains that comprise the alpha 5 subunit. This subunit associates with the beta 1 subunit to form a fibronectin receptor. This integrin may promote tumor invasion, and higher expression of this gene may be correlated with shorter survival time in lung cancer patients. Note that the integrin alpha 5 and integrin alpha V subunits are encoded by distinct genes.

# **Recommended Dilutions**

**WB** 1:500 - 1:1000

**ELISA** 

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

# Immunogen Information

**Gene ID**3678

Swiss Prot
P08648

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

# **Synonyms**

FNRA; CD49e; VLA-5; VLA5A; Integrin alpha 5 (ITGA5/CD49e)

# **Contact**

2		400-999-6126
$\bowtie$	Τ	cn.market@abclonal.com.cn
•	Т	www.abclonal.com.cn

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

# **Validation Data**



Western blot analysis of various lysates using Integrin alpha 5 (ITGA5/CD49e) Rabbit pAb (A22018) at 1:600 dilution

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3s.