

TriMethyl-Histone H3-K9 Rabbit mAb

Catalog No.: A22295

Recombinant

3 Publications

Basic Information

Observed MW

17kDa

Calculated MW

15kDa

Category

Primary antibody

Applications

WB, DB, IF/ICC, ELISA, ChIP, ChIP-seq, CUT&Tag

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

CloneNo number

ARC54898

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

WB 1:2000 - 1:20000**DB** 1:2000 - 1:20000**IF/ICC** 1:200 - 1:800**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.**ChIP** 5µg antibody for 5µg-10µg of Chromatin**ChIP-seq** 1:20 - 1:100**CUT&Tag** 10⁵ cells /1 µg

Immunogen Information

Gene ID

8290/8350

Swiss Prot

Q16695/P68431

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

H3.4; H3/g; H3FT; H3t; HIST3H3; Histone H3; HIST1H3A; TriMethyl-Histone H3-K9

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data




Western blot analysis of various lysates, using TriMethyl-Histone H3-K9 Rabbit mAb (A22295) at 1:20000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.




Chromatin immunoprecipitations were performed with cross-linked chromatin from HeLa cells and TriMethyl-Histone H3-K9 mAb (A22295). The ChIP sequencing results indicate the enrichment pattern of TriMethyl-Histone H3-K9 in selected genomic region and representative gene loci (ZNF404, GAPDH), as shown in figure.




Chromatin immunoprecipitation analysis of extracts of HeLa cells, using TriMethyl-Histone H3-K9 antibody (A22295) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Dot-blot analysis of all sorts of peptides using TriMethyl-Histone H3-K9 antibody (A22295) at 1:20000 dilution.



Confocal imaging of HeLa cells using TriMethyl-Histone H3-K9 Rabbit mAb (A22295, dilution 1:100) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) (Green). DAPI was used for nuclear staining (blue). Objective: HeLa.



CUT&Tag was performed using the CUT&Tag Assay Kit (pAG-Tn5) for Illumina (RK20265) from 10⁵ K562 cells with 1 µg TriMethyl-Histone H3-K9 Rabbit mAb (A22295), along with a Goat Anti-Rabbit IgG (H+L). The CUT&Tag results indicate the enrichment

Validation Data

100x.

pattern of TriMethyl-Histone H3-K9 in representative gene loci (MYT1), as shown in figure.