

[KO Validated] NF-κB p65/RelA Rabbit mAb

Catalog No.: A22331 **KO Validated** **Recombinant** **19 Publications**

Basic Information

Observed MW

65kDa

Calculated MW

58kDa/59kDa/60kDa

Category

Primary antibody

Applications

WB, IF/ICC, IP, ELISA, ChIP

Cross-Reactivity

Human, Mouse, Rat, Monkey

Clone/No. number

ARC51088

Background

NF-κB is a ubiquitous transcription factor involved in several biological processes. It is held in the cytoplasm in an inactive state by specific inhibitors. Upon degradation of the inhibitor, NF-κB moves to the nucleus and activates transcription of specific genes. NF-κB is composed of NFKB1 or NFKB2 bound to either REL, RELA, or RELB. The most abundant form of NF-κB is NFKB1 complexed with the product of this gene, RELA. Four transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB 1:5000 - 1:20000

IF/ICC 1:500 - 1:2000

IP 0.5μg-4μg antibody for 200μg-500μg extracts of whole cells

ChIP 5μg antibody for 10μg-15μg of Chromatin

ELISA Recommended starting concentration is 1 μg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

5970

Swiss Prot

Q04206

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

p65; CMCU; NFKB3; AIF3BL3; IA

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




Contact

 | 400-999-6126


 | cn.market@abclonal.com.cn






Validation Data



Validation Data



Chromatin immunoprecipitation analysis of extracts of HT-1080 cells, HT-1080 cells were treated by TNF- α (20 ng/ml) at 37°C for 30 minutes, using [KO Validated] NF-kB p65/RelA Rabbit mAb (A22331) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Confocal imaging of PC-12 cells (treated with TNF- α) and PC-12 cells (untreated) cells using [KO Validated] NF-kB p65/RelA Rabbit mAb (A22331, dilution 1:2000) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.

Confocal imaging of HT-1080 cells (treated with TNF- α) and HT-1080 cells (untreated) cells using [KO Validated] NF-kB p65/RelA Rabbit mAb (A22331, dilution 1:2000) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.

Confocal imaging of NIH/3T3 cells (treated with TNF- α) and NIH/3T3 cells (untreated) cells using [KO Validated] NF-kB p65/RelA Rabbit mAb (A22331, dilution 1:2000) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.