HDAC2 Rabbit mAb

Catalog No.: A22426 Recombinant 2 Publications



Basic Information

Observed MW

60kDa

Calculated MW

55kDa

Category

Primary antibody

Applications

WB,IHC-P,IF/ICC,IP,ELISA,ChIP

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC0105

Background

This gene product belongs to the histone deacetylase family. Histone deacetylases act via the formation of large multiprotein complexes, and are responsible for the deacetylation of lysine residues at the N-terminal regions of core histones (H2A, H2B, H3 and H4). This protein forms transcriptional repressor complexes by associating with many different proteins, including YY1, a mammalian zinc-finger transcription factor. Thus, it plays an important role in transcriptional regulation, cell cycle progression and developmental events. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:200 - 1:800

IF/ICC 1:50 - 1:200

IP 0.5μg-4μg antibody for

200μg-400μg extracts of

whole cells

ELISA Recommended starting

concentration is 1 µg/mL.

Please optimize the
concentration based on
your specific assay

requirements.

ELISA Recommended starting

concentration is 1 µg/mL.

Please optimize the
concentration based on
your specific assay
requirements.

ChIP 5μg antibody for

Immunogen Information

Gene IDSwiss Prot
3066
Q92769

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

HD2; RPD3; YAF1; KDAC2; HDAC2

Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


10μg-15μg of Chromatin.

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	Τ	www.abclonal.com.cn



Immunoprecipitation analysis of 300 μ g extracts from HeLa cells using 3 μ g HDAC2 Rabbit mAb (A22426). Western blot was performed from the immunoprecipitate using HDAC2 Rabbit mAb (A22426) at a dilution of 1:500.




Western blot analysis of lysates from HeLa cells, using HDAC2 Rabbit mAb (A22426) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 60s.




Immunofluorescence analysis of C6 cells using HDAC2 Rabbit mAb (A22426) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of NIH/3T3 cells using HDAC2 Rabbit mAb (A22426) at dilution of 1:100 (40xlens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of paraffinembedded Human colon carcinoma tissue using HDAC2 Rabbit mAb (A22426) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human liver cancer tissue using HDAC2 Rabbit mAb (A22426) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human lung squamous carcinoma tissue using HDAC2 Rabbit mAb (A22426) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffinembedded Mouse brain tissue using HDAC2 Rabbit mAb (A22426) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffinembedded Mouse intestin tissue using HDAC2 Rabbit mAb (A22426) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffinembedded Mouse liver tissue using HDAC2 Rabbit mAb (A22426) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Mouse spleen tissue using HDAC2 Rabbit mAb (A22426) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Rat brain tissue using HDAC2 Rabbit mAb (A22426) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Rat kidney tissue using HDAC2 Rabbit mAb (A22426) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.