

BCL6 Rabbit mAb

Catalog No.: A22610 **Recombinant**

Basic Information

Observed MW

Refer to figures

Calculated MW

79kDa

Category

Primary antibody

Applications

ELISA,IHC-P,FC (intra)

Cross-Reactivity

Human

CloneNo number

ARC57886

Background

The protein encoded by this gene is a zinc finger transcription factor and contains an N-terminal POZ domain. This protein acts as a sequence-specific repressor of transcription, and has been shown to modulate the transcription of STAT-dependent IL-4 responses of B cells. This protein can interact with a variety of POZ-containing proteins that function as transcription corepressors. This gene is found to be frequently translocated and hypermutated in diffuse large-cell lymphoma (DLCL), and may be involved in the pathogenesis of DLCL. Alternatively spliced transcript variants encoding different protein isoforms have been found for this gene.

Recommended Dilutions

IHC-P 1:100 - 1:500

FC (intra) 1:100 - 1:500

Immunogen Information

Gene ID

604

Swiss Prot

P41182

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 235-350 of human BCL6(NP_001124317.1).

Synonyms

BCL5; LAZ3; BCL6A; ZNF51; ZBTB27; BCL6

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.


Validation Data




Flow cytometry: 1×10^6 HAP1 cells (Low Expression, Left) and Daudi cells (Right) were intracellularly-stained with BCL6 Rabbit mAb (A22610, 2 $\mu\text{g/mL}$, orange line) or ABflo® 488 Rabbit IgG isotype control (A22069, 2 $\mu\text{g/mL}$, blue line), followed by FITC conjugated goat anti-rabbit pAb (1:200 dilution) staining. Non-fluorescently stained cells were used as blank control (red line).




Flow cytometry: 1×10^6 A20 cells were intracellularly-stained with BCL6 Rabbit mAb (A22610, 2 $\mu\text{g/mL}$, orange line) or ABflo® 488 Rabbit IgG isotype control (A22069, 2 $\mu\text{g/mL}$, blue line), followed by FITC conjugated goat anti-rabbit pAb (1:200 dilution) staining. Non-fluorescently stained A20 cells were used as blank control (red line).



Flow cytometry: 1×10^6 Daudi cells were intracellularly-stained with ABflo® 488 Rabbit IgG isotype control (A22069, 2 $\mu\text{g/mL}$, left) or BCL6 Rabbit mAb (A22610, 2 $\mu\text{g/mL}$, right).



Flow cytometry: 1×10^6 A20 cells were intracellularly-stained with ABflo® 488 Rabbit IgG isotype control (A22069, 2 $\mu\text{g/mL}$, left) or BCL6 Rabbit mAb (A22610, 2 $\mu\text{g/mL}$, right).



Immunohistochemistry analysis of BCL6 in paraffin-embedded Human tonsil using BCL6 Rabbit mAb (A22610) at dilution of 1:500 (40x lens). Perform high pressure antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.