

Uhrf2 Rabbit pAb

Catalog No.: A2344

1 Publications

Basic Information

Observed MW

95kDa

Calculated MW

90kDa

Category

Primary antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Enables histone binding activity. Predicted to be involved in several processes, including maintenance of DNA methylation; protein autoubiquitination; and regulation of cell cycle. Located in heterochromatin and nucleus. Is expressed in several structures, including central nervous system; ear; early conceptus; female reproductive system; and urinary system. Orthologous to human UHRF2 (ubiquitin like with PHD and ring finger domains 2).

Recommended Dilutions

WB 1:500 - 1:1000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

109113

Swiss Prot

Q7TMI3

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

Nirf; 2310065A22Rik; D130071B19Rik; Uhrf2

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG


Purification

Affinity purification

Storage

Store at -20°C

Validation Data



Western blot analysis of lysates from Mouse thymus, using Uhrf2 Rabbit pAb (A2344) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 180s.