Ų

www.abclonal.com

ABclonal

ABflo® 647 Rabbit anti-Human MMP9 mAb

Catalog No.: A23733

Basic Information

Observed MW

Calculated MW

78kDa

Category

Primary antibody

Applications

FC (intra)

Cross-Reactivity

Human

CloneNo number

ARC60377

Conjugate

ABflo® 647. Ex:648nm. Em:664nm.

Background

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The enzyme encoded by this gene degrades type IV and V collagens. Studies in rhesus monkeys suggest that the enzyme is involved in IL-8-induced mobilization of hematopoietic progenitor cells from bone marrow, and murine studies suggest a role in tumor-associated tissue remodeling.

Recommended Dilutions

FC (intra)

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID 4318 **Swiss Prot**

P14780

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

MMP9; CLG4B; GELB; MANDP2; MMP-9; matrix metallopeptidase 9

Contact

2		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Product Information

Source	Isotype	Purification
Rabbit	IgG	Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data









Flow cytometry: 1X10^6 293T cells(Low Expression,left) and 293T(Transfection,right) cells were intracellularly-stained with ABflo® 647 Rabbit anti-Human MMP9 mAb(A23733,5 µl/Test,orange line) or ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,blue line).Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:1X10^6 293T(Transfection) cells were intracellularly-stained with ABflo® 647 Rabbit IgG isotype control (A22070,5 μ I/Test,left) or ABflo® 647 Rabbit anti-Human MMP9 mAb(A23733,5 μ I/Test,right).