ABclonal www.abclonal.com

ABflo® 488 Rabbit anti-Human CD337/NKp30 mAb

Catalog No.: A23735

Basic Information

Observed MW

Calculated MW

16kDa/17kDa/18kDa/19kDa/20kDa/21kD

Category

Primary antibody

Applications

IF/ICC,FC (intra)

Cross-Reactivity

Human

CloneNo number

ARC60018

Conjugate

ABflo® 488. Ex:491nm. Em:516nm.

Background

The protein encoded by this gene is a natural cytotoxicity receptor (NCR) that may aid NK cells in the lysis of tumor cells. The encoded protein interacts with CD3-zeta (CD247), a T-cell receptor. A single nucleotide polymorphism in the 5' untranslated region of this gene has been associated with mild malaria suceptibility. Three transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

IF/ICC 1:50 - 1:200

FC (intra) 5 μl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID Swiss Prot 259197 014931

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

NCR3; 1C7; CD337; LY117; MALS; NKp30; natural cytotoxicity triggering receptor 3

Contact

a	400-999-6126
\bowtie	cn.market@abclonal.com.cn
<u></u>	www.abclonal.com.cn

Product Information


Source	Isotype	Purification
Rabbit	IgG	Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data










Flow cytometry: 1X10^6 293T cells(Low Expression,left) and 293T(Transfection,right) cells were intracellularly-stained with ABflo® 488 Rabbit anti-Human CD337/NKp30 mAb(A23735,5 µl/Test,orange line) or ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,blue line).Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:1X10^6 293T(Transfection) cells were intracellularly-stained with ABflo® 488 Rabbit IgG isotype control (A22069,5 μ I/Test,left) or ABflo® 488 Rabbit anti-Human CD337/NKp30 mAb(A23735,5 μ I/Test,right).



Confocal imaging of 293T cells transfected with CD337/NKp30 using ABflo® 488 Rabbit anti-Human CD337/NKp30 mAb (A23735, dilution 1:200). DAPI was used for nuclear staining (Blue). Objective: 100x.