

VEGF Rabbit mAb

Catalog No.: A23759

Recombinant

14 Publications

Basic Information

Observed MW

16kDa/35kDa

Calculated MW

15-27kDa/34-45kDa

Category

Primary antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC61370

Background

This gene is a member of the PDGF/VEGF growth factor family. It encodes a heparin-binding protein, which exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascular endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Elevated levels of this protein are found in patients with POEMS syndrome, also known as Crow-Fukase syndrome. Allelic variants of this gene have been associated with microvascular complications of diabetes 1 (MVCD1) and atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been described. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in additional isoforms. A recent study showed that a C-terminally extended isoform is produced by use of an alternative in-frame translation termination codon via a stop codon readthrough mechanism, and that this isoform is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a small upstream open reading frame, which is located within an internal ribosome entry site.

Recommended Dilutions

| | |
|-----------|------------------|
| WB | 1:3000 - 1:12000 |
|-----------|------------------|

| | |
|---------------|---------------|
| IF/ICC | 1:200 - 1:800 |
|---------------|---------------|

| | |
|--------------|---|
| ELISA | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |
|--------------|---|

Immunogen Information

Gene ID

7422

Swiss Prot

P15692

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

VEGFA; MVCD1; VEGF; VPF; vascular endothelial growth factor A

Contact

| | | |
|--|--|--------------|
|  | | 400-999-6126 |
|--|--|--------------|

| | | |
|--|--|---------------------------|
|  | | cn.market@abclonal.com.cn |
|--|--|---------------------------|

| | | |
|--|--|---------------------|
|  | | www.abclonal.com.cn |
|--|--|---------------------|

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data




Western blot analysis of recombinant Human VEGF-A/VEGF121(K321N) Protein using VEGF Rabbit mAb (A23759) at 1:3000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 10ng per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.



Immunofluorescence analysis of C6 cells using VEGF Rabbit mAb (A23759) at a dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of K-562 cells using VEGF Rabbit mAb (A23759) at a dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using VEGF Rabbit mAb (A23759) at a dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.