Leader in Biomolecular Solutions for Life Science

www.abclonal.com

ABflo® 594 Rabbit anti-Mouse/Human CD55

Catalog No.: A23899

Basic Information

Observed MW

Calculated MW
42KDa

Category
Primary antibody

Applications
FC

Cross-Reactivity
Human, Mouse

CloneNo number
ARC60934

Conjugate
ABflo® 594. Ex:588nm. Em:604nm.

Background

This gene encodes an inhibitor of both the classical and the alternative pathways of complement activation. The encoded preproprotein undergoes post-translational processing to generate a mature polypeptide anchored to the plasma membrane via a glycosylphosphatidylinositol moiety. Erythrocytes from mice deficient in the encoded protein exhibit impaired regulation of complement activation resulting in enhanced complement deposition. Mice lacking the encoded protein exhibit enhanced susceptibility to experimentally induced myasthenia gravis. This gene is located adjacent to a closely related gene on chromosome 1.

Recommended Dilutions

FC	5 µl per 10^6 cells in 100 µl volume

Immunogen Information

Gene ID
13136

Swiss Prot
Q61475

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
Daf; Daf1; Daf-GPI; GPI-DAF

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Product Information

Source
Rabbit

Isotype
IgG

Purification
Affinity purification

Storage
Store at -20℃

Validation Data



Flow cytometry: 1X10^6 C57BL/6 mouse Splenocytes were surface-stained with ABflo® 594 Rabbit anti-Human/Mouse CD55/DAF mAb(A23899,5 µl/Test,orange line) or ABflo® 594 Rabbit IgG isotype control (A23821,5 µl/Test,blue line).Non-fluorescently stained C57BL/6 mouse Splenocytes were used as blank control (red line).



Flow cytometry: 1X10^6 SH-SY5Y cells(negative control,left) and Hela cells(right) were surface-stained with ABflo® 594 Rabbit anti-Human/Mouse CD55/DAF mAb(A23899,5 µl/Test,orange line) or ABflo® 594 Rabbit IgG isotype control (A23821,5 µl/Test,blue line).Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry:1X10^6 C57BL/6 mouse Splenocytes were surface-stained with ABflo® 594 Rabbit IgG isotype control (A23821,5 µl/Test,left) or ABflo® 594 Rabbit anti-Human/Mouse CD55/DAF mAb(A23899,5 µl/Test,right).



Flow cytometry:1X10^6 Hela cells were surface-stained with ABflo® 594 Rabbit IgG isotype control (A23821,5 µl/Test,left) or ABflo® 594 Rabbit anti-Human/Mouse CD55/DAF mAb(A23899,5 µl/Test,right).