

c-Fos Rabbit mAb

Catalog No.: A24620 **Recombinant** **3 Publications**

Basic Information

Observed MW

62 kDa

Calculated MW

41 kDa

Category

Primary antibody

Applications

WB,IF-P,IHC-P,mIHC,ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC63309

Background

The Fos gene family consists of 4 members: FOS, FOSB, FOSL1, and FOSL2. These genes encode leucine zipper proteins that can dimerize with proteins of the JUN family, thereby forming the transcription factor complex AP-1. As such, the FOS proteins have been implicated as regulators of cell proliferation, differentiation, and transformation. In some cases, expression of the FOS gene has also been associated with apoptotic cell death.

Recommended Dilutions

WB	1:1000 - 1:4000
IF-P	1:200 - 1:400
IHC-P	1:200 - 1:800
mIHC	1:500 - 1:2000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Immunogen Information

Gene ID

2353

Swiss Prot

P01100

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

p55; AP-1; C-FOS; c-Fos

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data



Western blot analysis of lysates from NIH/3T3 cells using c-Fos Rabbit mAb (A24620) at 1:1000 dilution incubated overnight at 4°C. NIH/3T3 cells were treated with PMA(200 nM) at 37°C for 30 minutes after serum-starvation overnight.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 30 µg per lane.
Blocking buffer: 3 % nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 45s.



Western blot analysis of lysates from PC-12 cells using c-Fos Rabbit mAb (A24620) at 1:1000 dilution incubated overnight at 4°C. PC-12 cells were treated with PMA(200 nM) at 37°C for 30 minutes after serum-starvation overnight.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 30 µg per lane.
Blocking buffer: 3 % nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 60s.




Western blot analysis of lysates from HeLa cells using c-Fos Rabbit mAb (A24620) at 1:1000 dilution incubated overnight at 4°C. HeLa cells were treated with PMA(200 nM) at 37°C for 30 minutes after serum-starvation overnight.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3 % nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 45s.


Validation Data




Immunohistochemistry analysis of paraffin-embedded Human cervix using c-Fos Rabbit mAb (A24620) at dilution of 1:400 (40x lens). High pressure antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human colon using c-Fos Rabbit mAb (A24620) at dilution of 1:400 (40x lens). High pressure antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.




Confocal imaging of paraffin-embedded Mouse brain tissue using c-Fos Rabbit mAb (A24620, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of paraffin-embedded Mouse colon tissue using c-Fos Rabbit mAb (A24620, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




Confocal imaging of paraffin-embedded Rat brain tissue using c-Fos Rabbit mAb (A24620, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.




The multiplex IHC analysis on paraffin-embedded Rat brain tissue using the following specific primary antibodies and tyramide signal amplification (TSA) reagents (RK05903) : c-Fos Rabbit mAb (A24620, 1:1000) with TSA-TYR-520 (Green), and GAD67/GAD1 Rabbit mAb (A26748, 1:500) with TSA-TYR-570 (Red). DAPI (Blue) was used for nuclear staining. Prior to multiplex IHC staining, high-pressure antigen retrieval was performed using 0.01M citrate buffer at pH 6.0. The analysis was completed using a 40x objective lens.



The multiplex IHC analysis on paraffin-embedded Mouse brain tissue using the



3D imaging of solvent-cleared Mouse brain tissue using c-Fos Rabbit mAb (A24620,



3D imaging of solvent-cleared Mouse brain tissue using c-Fos Rabbit mAb (A24620,

Validation Data

following specific primary antibodies and tyramide signal amplification (TSA) reagents (RK05903) : c-Fos Rabbit mAb (A24620, 1:1000) with TSA-TYR-520 (Green), and GAD67/GAD1 Rabbit mAb (A26748, 1:500) with TSA-TYR-570 (Red). DAPI (Blue) was used for nuclear staining. Prior to multiplex IHC staining, high-pressure antigen retrieval was performed using 0.01M citrate buffer at pH 6.0. The analysis was completed using a 40x objective lens.

dilution 1:200). SailClear Tissue Optical Clearing Kit(QF2601) was used for sample clearing. We acknowledge SailLife(Nanjing) Sci-Tech Co., Ltd. in 3D imaging processing and kindly providing this image.

dilution 1:200). SailClear Tissue Optical Clearing Kit(QF2601) was used for sample clearing. We acknowledge SailLife(Nanjing) Sci-Tech Co., Ltd. in 3D imaging processing and kindly providing this image.