

ABflo® 647 Rabbit anti-Mouse CD127/IL-7R α mAb

Catalog No.: A24789

Basic Information

Observed MW

Calculated MW

51kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC63550

Conjugate

ABflo® 647. Ex:648nm. Em:664nm.

Background

Interleukin-7 is a glycoprotein involved in the regulation of lymphopoiesis. Response of cells to IL7 is dependent on the presence of the interleukin 7 receptor (IL7R); the active receptor is a α/γ chain heterodimer. The $\gamma(c)$ chain, which also associates with the interleukin-2 receptor, serves primarily to activate signal transduction by the IL7R complex, while the α chain of IL7R determines specific signaling events through its association with cytoplasmic signaling molecules.

Recommended Dilutions

FC 5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID

16197

Swiss Prot

P16872

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CD127; IL-7R α

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Flow cytometry: 1×10^6 C57BL/6 mouse splenocytes were surface-stained with ABflo® 488 Rat anti-Mouse CD3 mAb (A27161, 5 μ l/Test) and ABflo® 647 Rabbit IgG isotype control (A22070, 5 μ l/Test, left) or ABflo® 647 Rabbit anti-Mouse CD127/IL-7R α mAb (A24789, 5 μ l/Test, right).