# PRELID1 Rabbit pAb

Catalog No.: A25108



### **Basic Information**

#### **Observed MW**

Refer to figures

### **Calculated MW**

25kDa

### Category

Primary antibody

### **Applications**

IHC-P,IF/ICC,ELISA

#### **Cross-Reactivity**

Human

# **Background**

This gene encodes a member of the late embryogenesis abundant motif-containing protein family. The encoded protein is localized to mitochondria and may function as a cytoprotectant by regulating cell death and differentiation. Alternative splicing results in multiple transcript variants encoding different isoforms. Several related pseudogenes have been identified.

## **Recommended Dilutions**

**IHC-P** 1:50 - 1:200

**IF/ICC** 1:50 - 1:200

Recommended starting concentration is 1 µg/mL.

Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**27166

Swiss Prot
29Y255

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

PX19; PRELI; SBBI12; CGI-106

### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	Ī	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification


### **Storage**

Store at -20  $^{\circ}\text{C}.$  Avoid freeze / thaw cycles.


Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Immunofluorescence analysis of Hep G2 cells using PRELID1 Rabbit pAb(A25108) at a dilution of 1:100 (40x lens). Secondary antibody:Cy3 Goat Anti-Rabbit IgG (H+L)(AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffinembedded Human colon carcinoma tissue using PRELID1 Rabbit pAb (A25108) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Human tonsil tissue using PRELID1 Rabbit pAb (A25108) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.