

P-cadherin Rabbit mAb

Catalog No.: A25549 **Recombinant**

Basic Information

Observed MW

120kDa

Calculated MW

87kDa/91kDa

Category

Primary antibody

Applications

WB,IF/ICC,IF-P,FC,ELISA

Cross-Reactivity

Human

CloneNo number

ARC66488

Background

This gene encodes a classical cadherin of the cadherin superfamily. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate the mature glycoprotein. This calcium-dependent cell-cell adhesion protein is comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. This gene is located in a gene cluster in a region on the long arm of chromosome 16 that is involved in loss of heterozygosity events in breast and prostate cancer. In addition, aberrant expression of this protein is observed in cervical adenocarcinomas. Mutations in this gene are associated with hypotrichosis with juvenile macular dystrophy and ectodermal dysplasia, ectrodactyly, and macular dystrophy syndrome (EEMS).


Recommended Dilutions

WB	1:3000 - 1:12000
IF/ICC	1:200 - 1:800
IF-P	1:200 - 1:800
FC	1:500 - 1:1000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Immunogen Information

Gene ID

1001

Swiss Prot

P22223

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CDHP; HJMD; PCAD

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data




Western blot analysis of various lysates using P-cadherin Rabbit mAb (A25549) at 1:3000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Negative control (NC): SH-SY5Y. Exposure time: 30s.




Flow cytometry: 1×10^6 MCF7 cells (Low Expression, left) and A-431 cells (right) were surface-stained with P-cadherin Rabbit mAb (A25549, 2 µg/mL, orange line) or ABflo® 647 Rabbit IgG isotype control (A22070, 5 µg/mL, blue line), followed by Alexa Fluor® 647 conjugated goat anti-rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1×10^6 A-431 cells were surface-stained with ABflo® 647 Rabbit IgG isotype control (A22070, 5 µg/mL, left) or P-cadherin Rabbit mAb (A25549, 2 µg/mL, right).



Confocal immunofluorescence analysis of A-431 cells using P-cadherin Rabbit mAb (A25549, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Confocal imaging of paraffin-embedded Human breast cancer tissue using P-cadherin Rabbit mAb (A25549, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval with 0.01 M citrate buffer (pH 6.0) prior to IF staining. Objective: 40x.