RACK1 Rabbit pAb

Catalog No.: A2560 1 Publications



Basic Information

Observed MW

32kDa

Calculated MW

35kDa

Category

Primary antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Enables several functions, including cyclin binding activity; enzyme binding activity; and protein domain specific binding activity. Involved in several processes, including positive regulation of hydrolase activity; regulation of cellular protein metabolic process; and regulation of signal transduction. Located in several cellular components, including midbody; perinuclear region of cytoplasm; and phagocytic cup. Part of IRE1-RACK1-PP2A complex.

Recommended Dilutions

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:200

ELISA Recommended starting concentration is 1 μg/mL.

Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID Swiss Prot 10399 P63244

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

H12.3; HLC-7; PIG21; GNB2L1; Gnb2-rs1; RACK1

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\odot	T	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of various lysates using RACK1 Rabbit pAb (A2560) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: $25\mu g$ per lane.

Blocking buffer: 3% nonfat dry milk in TBST.



Western blot analysis of various lysates using RACK1 Rabbit pAb (A2560) at 1:1000 dilution incubated overnight at 4° C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 μg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 10s.



Western blot analysis of various lysates using RACK1 Rabbit pAb (A2560) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: $25\mu g$ per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Validation Data



Immunofluorescence analysis of MCF-7 cells using RACK1 Rabbit pAb (A2560).Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution.