FITC anti-Mouse CD45R/B220 mAb

Catalog No.: A25655



Basic Information

Observed MW

Refer to figures

Calculated MW

180-240kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ATC0011

Conjugate

FITC. Ex:491nm. Em:516nm.

Background

Enables several functions, including heparan sulfate proteoglycan binding activity; heparin binding activity; and protein tyrosine phosphatase activity. Involved in several processes, including lymphocyte differentiation; positive regulation of macromolecule metabolic process; and regulation of signal transduction. Acts upstream of or within several processes, including lymphocyte differentiation; positive regulation of lymphocyte activation; and regulation of protein phosphorylation. Located in external side of plasma membrane; focal adhesion; and membrane raft. Is expressed in several structures, including 3rd branchial arch; alimentary system; cardiovascular system; hemolymphoid system; and placenta. Used to study systemic lupus erythematosus. Human ortholog(s) of this gene implicated in hepatitis C; multiple sclerosis; severe combined immunodeficiency; and severe combined immunodeficiency, autosomal recessive, T cell-negative, B cell-positive, Nk cell-positive. Orthologous to human PTPRC (protein tyrosine phosphatase receptor type C).

Recommended Dilutions

FC

The suggested use of this reagent is ≤1.0 µg per 10^6 cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application

Immunogen Information

Gene ID 19264

Swiss Prot P08575/P06800

Immunogen

This information is considered to be commercially sensitive.

Synonyms

loc; B220; Cd45; L-CA; Ly-5; T200; CD45R; Lyt-4

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\odot	T	www.abclonal.com.cn

Product Information

SourceIsotypePurificationRatRat IgG2a, κAffinity purification

Storage

Store at 4°C. Avoid freeze / thaw cycles. Buffer:Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide