IL12 Rabbit pAb

Catalog No.: A25798



Basic Information

Observed MW

40-55kDa

Calculated MW

25kDa/37kDa

Category

Primary antibody

Applications

ELISA,WB

Cross-Reactivity

Human

Background

This gene encodes a subunit of a cytokine that acts on T and natural killer cells, and has a broad array of biological activities. The cytokine is a disulfide-linked heterodimer composed of the 35-kD subunit encoded by this gene, and a 40-kD subunit that is a member of the cytokine receptor family. This cytokine is required for the T-cell-independent induction of interferon (IFN)-gamma, and is important for the differentiation of both Th1 and Th2 cells. The responses of lymphocytes to this cytokine are mediated by the activator of transcription protein STAT4. Nitric oxide synthase 2A (NOS2A/NOS2) is found to be required for the signaling process of this cytokine in innate immunity.

Recommended Dilutions

WB

1:500 - 1:1000

Immunogen Information

Gene ID 3592/3593

Swiss Prot P29459/P29460

_

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 23-328 of human IL12B/23-219 of human IL12A (NP_002178.2/NP_001384921.1).

Synonyms

P35; CLMF; NFSK; NKSF1; IL-12A

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	ī	www.abclonal.com.cn


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of lysates from recombinant Human IL-12B/P40 Protein cells using IL12 Rabbit pAb (A25798) at 1:1000 dilution incubated overnight at 4°C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L)(AS014) at 1:10000 dilution.

Lysates/proteins: 0.5-5ng μg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020) Exposure time: 90 s.