ABclonal www.abclonal.com

ABflo® 610 Rabbit IgG isotype control

Catalog No.: A25826

Basic Information

Observed MW

Calculated MW

Category

Primary antibody

Applications

FC

Cross-Reactivity

CloneNo number

ARC5105-10-ABf610

Conjugate

ABflo® 610. Ex:421nm. Em:612nm.

Background

The isotype of a primary antibody and the application it is being used in can result in background staining. Primary antibody background noise can be caused by binding to Fc receptors on target cells; by non-specific interactions with cellular proteins, carbohydrates, and lipids; or by cell autofluorescence. Isotype control antibodies can act as negative controls to help differentiate non-specific background signal from specific antibody signal because they have no relevant specificity to a target antigen. An isotype control antibody should have the same immunoglobulin type and be used at the same concentration as the test antibody.

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID

Swiss Prot

Immunogen

A synthesized peptide derived from rabbit IgG isotype control.

Synonyms

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
\odot	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

Validation Data





Flow cytometry: 1X10^6 293F cells (negative control,left) and Human PBMC (right) were surface-stained with ABflo® 610 Rabbit anti-Human CD8a mAb (A25765,5 µl/Test,orange line) or ABflo® 610 Rabbit IgG isotype control (A25826,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: $1X10^6$ Human PBMC were surface-stained with ABflo® 610 Rabbit IgG isotype control (A25826,5 μ I/Test,left) or ABflo® 610 Rabbit anti-Human CD8a mAb (A25765,5 μ I/Test,right).