CD172b Rabbit mAb

Catalog No.: A25883 Recombinant



Basic Information

Observed MW

Calculated MW 19kDa/43kDa

Category

Primary antibody

Applications

ELISA,FC

Cross-Reactivity

Human

CloneNo number

ARC67299

Background

The protein encoded by this gene is a member of the signal-regulatory-protein (SIRP) family, and also belongs to the immunoglobulin superfamily. SIRP family members are receptor-type transmembrane glycoproteins known to be involved in the negative regulation of receptor tyrosine kinase-coupled signaling processes. This protein was found to interact with TYROBP/DAP12, a protein bearing immunoreceptor tyrosine-based activation motifs. This protein was also reported to participate in the recruitment of tyrosine kinase SYK. Multiple transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

FC

1:500 - 1:1000

Immunogen Information

Gene ID 10326

Swiss Prot

000241

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 30-371 of human CD172b (NP_006056.2).

Synonyms

CD172b; SIRP-BETA-1

Contact

6		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20 $^{\circ}\text{C}.$ Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

Validation Data







Flow cytometry: 1X10^6 HEL cells (negative control,left) and U-937 cells (right) cells were surface-stained with CD172b Rabbit mAb (A25883,2 μg/mL,orange line) or Rabbit lgG isotype control (AC042,2 μg/mL,blue line), followed by Alexa Fluor® 647 conjugated goat anti-rabbit pAb staining. Nonfluorescently stained cells were used as blank control (red line).

Flow cytometry: 1X10^6 U-937 cells were surface-stained with Rabbit IgG isotype control (AC042,2 µg/mL,left) or CD172b Rabbit mAb (A25883,2 µg/mL,right), followed by Alexa Fluor® 647 conjugated goat antirabbit pAb staining.