

# ABflo® 700 Rabbit IgG isotype control

Catalog No.: A25976

## Basic Information

---

### Observed MW

### Calculated MW

### Category

Primary antibody

### Applications

FC

### Cross-Reactivity

### CloneNo number

ARC5105-10

### Conjugate

ABflo® 700. Ex:690nm. Em:713nm.

## Background

---

The isotype of a primary antibody and the application it is being used in can result in background staining. Primary antibody background noise can be caused by binding to Fc receptors on target cells; by non-specific interactions with cellular proteins, carbohydrates, and lipids; or by cell autofluorescence. Isotype control antibodies can act as negative controls to help differentiate non-specific background signal from specific antibody signal because they have no relevant specificity to a target antigen. An isotype control antibody should have the same immunoglobulin type and be used at the same concentration as the test antibody.

## Recommended Dilutions

---

FC 5 µl per 10<sup>6</sup> cells in  
100 µl volume

## Immunogen Information

---

### Gene ID

### Swiss Prot

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

## Contact

---

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

---

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Validation Data



Flow cytometry:  $1 \times 10^6$  293T cells (negative control, left) and Jurkat cells (right) were surface-stained with ABflo® 700 Rabbit anti-Human CD45 mAb ( $2 \mu\text{g}/\text{mL}$ , orange line) or ABflo® 700 Rabbit IgG isotype control ( $2.5 \mu\text{g}/\text{mL}$ , blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:  $1 \times 10^6$  Jurkat cells were surface-stained with ABflo® 700 Rabbit IgG isotype control ( $2.5 \mu\text{g}/\text{mL}$ , left) or ABflo® 700 Rabbit anti-Human CD45 mAb ( $2 \mu\text{g}/\text{mL}$ , right).