

APC Rabbit anti-Mouse CD96/TACTILE mAb

Catalog No.: A26001

Basic Information

Observed MW

Calculated MW

67kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC63717

Conjugate

APC. Ex:650nm. Em:660nm.

Recommended Dilutions

FC 5 μ l per 10^6 cells in
100 μ l volume

Background

Acts upstream of or within negative regulation of interferon-gamma production; negative regulation of natural killer cell cytokine production; and response to lipopolysaccharide. Predicted to be located in cytoplasm. Is expressed in several structures, including blood; central nervous system; ganglia; heart; and lung. Human ortholog(s) of this gene implicated in C syndrome. Orthologous to human CD96 (CD96 molecule).

Immunogen Information

Gene ID

84544

Swiss Prot

Q3U0X8

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

Tactile; 1700109112Rik

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification



Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1×10^6 C2C12 cells (negative control, left) and C57BL/6 Mouse Splenocytes (right) were surface-stained with APC Rabbit anti-Mouse CD96/TACTILE mAb ($2.6 \mu\text{l}/\text{Test}$, orange line) or APC Rabbit IgG isotype control ($2.4 \mu\text{l}/\text{Test}$, blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1×10^6 C57BL/6 mouse splenocytes were surface-stained with APC Rabbit IgG isotype control ($2.4 \mu\text{l}/\text{Test}$, left) or APC Rabbit anti-Mouse CD96/TACTILE mAb ($2.6 \mu\text{l}/\text{Test}$, right).