

PE Rabbit anti-4-Hydroxynonenal/4-HNE mAb

Catalog No.: A26084

Basic Information

Observed MW

Calculated MW

Category

Primary antibody

Applications

FC (intra)

Cross-Reactivity

Species independent

CloneNo number

ARC70314

Conjugate

PE. Ex:565nm. Em:574nm.

Background

4-hydroxy-2-nonenal (4-hydroxynonenal, 4-HNE) is a highly reactive aldehyde generated by the exposure of polyunsaturated fatty acids to peroxides and reactive oxygen species (ROS). It non-enzymatically forms stable protein adducts with histidine, lysine, and cysteine side chains that have been used as biomarkers for oxidative damage in cells. Conditions where 4-HNE immunoreactivity has been observed include inflammation, neurodegenerative diseases, and ischemic damage to the heart and brain.

Recommended Dilutions

FC (intra) 5 μ l per 10^6 cells in
100 μ l volume

Immunogen Information

Gene ID

Swiss Prot

Immunogen

Chemical compounds corresponding to 4-Hydroxynonenal.

Synonyms

4-HNE

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1×10^6 HeLa cells (treated with 4-Hydroxynonenal) were intracellularly-stained with PE Rabbit IgG isotype control (A24172, 5 μ l/Test, left) or PE Rabbit anti-4-Hydroxynonenal mAb (A26084, 5 μ l/Test, right).