

APC Rabbit anti-Human CD40 mAb

Catalog No.: A26130

Basic Information

Observed MW**Calculated MW**

31kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC57632

Conjugate

APC. Ex:650nm. Em:660nm.

Background

This gene is a member of the TNF-receptor superfamily. The encoded protein is a receptor on antigen-presenting cells of the immune system and is essential for mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation. AT-hook transcription factor AKNA is reported to coordinately regulate the expression of this receptor and its ligand, which may be important for homotypic cell interactions. Adaptor protein TNFR2 interacts with this receptor and serves as a mediator of the signal transduction. The interaction of this receptor and its ligand is found to be necessary for amyloid-beta-induced microglial activation, and thus is thought to be an early event in Alzheimer disease pathogenesis. Mutations affecting this gene are the cause of autosomal recessive hyper-IgM immunodeficiency type 3 (HIGM3). Multiple alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.

Recommended Dilutions

FC 5 µl per 10⁶ cells in
100 µl volume

Immunogen Information

Gene ID

958

Swiss Prot

P25942

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

p50; Bp50; CDW40; TNFRSF5

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.


Validation Data




Flow cytometry: 1×10^6 Jurkat cells (negative control, left) and U-2 OS cells (right) were surface-stained with APC Rabbit anti-Human CD40 mAb (A26130, 5 µl/Test, orange line) or APC Rabbit IgG isotype control (A24173, 5 µl/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1×10^6 U-2 OS cells were surface-stained with APC Rabbit IgG isotype control (A24173, 5 µl/Test, left) or APC Rabbit anti-Human CD40 mAb (A26130, 5 µl/Test, right).



Flow cytometry: 1×10^6 Human PBMC were surface-stained with APC Rabbit anti-Human CD40 mAb (A26130, 5 µl/Test, orange line) or APC Rabbit IgG isotype control (A24173, 5 µl/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1×10^6 Human PBMC were surface-stained with APC Rabbit IgG isotype control (A24173, 5 µl/Test, left) or APC Rabbit anti-Human CD40 mAb (A26130, 5 µl/Test, right).