

ABflo® 647 Rabbit anti-Mouse E-Cadherin/CD324 mAb

Catalog No.: A26484

Basic Information

Observed MW

Calculated MW

98kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC68219

Conjugate

ABflo® 647. Ex:648nm. Em:664nm.

Background

This gene encodes E-cadherin, a calcium-dependent cell adhesion molecule that functions in the establishment and maintenance of epithelial cell morphology during embryongenesis and adulthood. The encoded preproprotein undergoes proteolytic processing to generate a mature protein. Targeted mutations disrupting binding of calcium to the encoded protein in mice cause death in utero due to failed blastocyst and trophectoderm formation. This gene is located adjacent to a related cadherin gene on chromosome 8.

Recommended Dilutions

FC

5 μl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID 12550

Swiss Prot

P09803

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

Um; UVO; Ecad; ARC-1; E-cad; L-CAM

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\odot	Т	www.abclonal.com.cn

Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data









Flow cytometry: 1X10^6 CHO-S cells (negative control,left) and CHO-S (Transfection,right) cells were surface-stained with ABflo® 647 Rabbit anti-Mouse E-Cadherin/CD324 mAb (A26484,5 µl/Test,orange line) or ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1X10^6 CHO-S (Transfection) cells were surface-stained with ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,left) or ABflo® 647 Rabbit anti-Mouse E-Cadherin/CD324 mAb (A26484,5 µl/Test,right).