APC Rabbit anti-Human CD11a mAb

www.abclonal.com

ABclonal

Catalog No.: A26541

Basic Information

Observed MW

Calculated MW

118kDa/129kDa/134kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC62732

Conjugate

APC. Ex:650nm. Em:660nm.

Background

ITGAL encodes the integrin alpha L chain. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This I-domain containing alpha integrin combines with the beta 2 chain (ITGB2) to form the integrin lymphocyte function-associated antigen-1 (LFA-1), which is expressed on all leukocytes. LFA-1 plays a central role in leukocyte intercellular adhesion through interactions with its ligands, ICAMs 1-3 (intercellular adhesion molecules 1 through 3), and also functions in lymphocyte costimulatory signaling. Two transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID 3683 Swiss Prot

P20701

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CD11A; LFA-1; LFA1A

Contact

a	400-999-6126
\sim	cn.market@abclonal.com.cn
⊙	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data





Flow cytometry: 1X10^6 K-562 cells (negative control,left) and Human PBMC (right) were surface-stained with APC Rabbit anti-Human CD11a mAb (A26541,5 μ l/Test,orange line) or APC Rabbit IgG isotype control (A24173,5 μ l/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: $1X10^6$ Human PBMC were surface-stained with APC Rabbit IgG isotype control (A24173,5 μ I/Test,Ieft) or APC Rabbit anti-Human CD11a mAb (A26541,5 μ I/Test,right).