# PE Rabbit anti-Human mGluR2 mAb

www.abclonal.com

**ABclonal** 

Catalog No.: A27130

### **Basic Information**

#### **Observed MW**

### **Calculated MW**

96kDa

### Category

Primary antibody

## **Applications**

FC

#### **Cross-Reactivity**

Human

#### CloneNo number

ARC70955

### Conjugate

PE. Ex:565nm. Em:574nm.

## **Background**

L-glutamate is the major excitatory neurotransmitter in the central nervous system and activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors, that have been divided into 3 groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Several transcript variants encoding different isoforms have been found for this gene.

### **Recommended Dilutions**

FC

5 μl per 10^6 cells in 100 μl volume

## Immunogen Information

Gene ID 2912 **Swiss Prot** 

Q14416

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

GLUR2; mGlu2; GPRC1B; MGLUR2

### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	T	www.abclonal.com.cn

### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

## **Validation Data**









Flow cytometry: 1X10^6 293T cells (negative control,left) and 293T (Transfection,right) cells were surface-stained with PE Rabbit anti-Human mGluR2 mAb (A27130,5 µl/Test,orange line) or PE Rabbit IgG isotype control (A24172,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:  $1X10^6$  293T (Transfection) cells were surface-stained with PE Rabbit IgG isotype control (A24172,5  $\mu$ I/Test,Ieft) or PE Rabbit anti-Human mGIuR2 mAb (A27130,5  $\mu$ I/Test,right).