ABflo® 647 Rabbit anti-Mouse CD40 mAb

ABclonal www.abclonal.com

Catalog No.: A27176

Basic Information

Observed MW

Calculated MW

32kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC71300

Conjugate

ABflo® 647. Ex:648nm. Em:664nm.

Background

Predicted to enable antigen binding activity; protein domain specific binding activity; and ubiquitin protein ligase binding activity. Involved in B cell mediated immunity; CD40 signaling pathway; and cellular calcium ion homeostasis. Acts upstream of or within several processes, including defense response to other organism; positive regulation of B cell activation; and positive regulation of interleukin-12 production. Located in external side of plasma membrane and intracellular membrane-bounded organelle. Part of CD40 receptor complex. Is expressed in several structures, including alimentary system; brain; hemolymphoid system gland; liver and biliary system; and reproductive system. Human ortholog(s) of this gene implicated in several diseases, including Kawasaki disease; autoimmune disease (multiple); end stage renal disease; hyperimmunoglobulin syndrome (multiple); and non-Hodgkin lymphoma (multiple). Orthologous to human CD40 (CD40 molecule).

Recommended Dilutions

FC

5 μl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID 21939

Swiss Prot

P27512

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

IGM; p50; Bp50; GP39; IMD3; TRAP; HIGM1; T-BAM; Tnfrsf5

Contact

| a | | 400-999-6126 |
|-----------|---|---------------------------|
| \bowtie | | cn.market@abclonal.com.cn |
| \odot | T | www.abclonal.com.cn |


Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.





Flow cytometry: 1X10^6 C57BL/6 mouse splenocytes were surface-stained with ABflo® 647 Rabbit anti-Mouse CD40 mAb (A27176,5 µl/Test,orange line) or ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line). Cells in the gate were used for analysis.

Flow cytometry: 1X10^6 C57BL/6 mouse splenocytes were surface-stained with ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,left) or ABflo® 647 Rabbit anti-Mouse CD40 mAb (A27176,5 µl/Test,right).