# CYP39A1 Rabbit mAb

Catalog No.: A27264 Recombinant



## **Basic Information**

#### **Observed MW**

55kDa

### **Calculated MW**

54kDa

### Category

Primary antibody

### **Applications**

WB,IHC-P,ELISA

### **Cross-Reactivity**

Mouse

#### CloneNo number

ARC69091

## **Background**

This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This endoplasmic reticulum protein is involved in the conversion of cholesterol to bile acids. Its substrates include the oxysterols 25-hydroxycholesterol, 27-hydroxycholesterol and 24-hydroxycholesterol. Alternate splicing results in multiple transcript variants.

# **Recommended Dilutions**

**WB** 1:14000 - 1:56000

IHC-P 1:2000 - 1:7000

**ELISA** Recommended starting concentration is 1 μg/mL.

Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**Swiss Prot
51302
Q9NYL5

### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

**Synonyms** 

### **Contact**

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\overline{a}$	ī	www.ahclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Western blot analysis of lysates from Mouse liver using CYP39A1 Rabbit mAb (A27264) at 1:14000 dilution incubated overnight at  $4^{\circ}$ C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 20s.



Immunohistochemistry analysis of paraffinembedded Mouse liver tissue using CYP39A1 Rabbit mAb (A27264) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.