

PE/Cyanine7 Rabbit anti-Mouse CD11b mAb

Catalog No.: A27339

Basic Information

Observed MW

Calculated MW
127kDa

Category
Primary antibody

Applications
FC

Cross-Reactivity
Mouse

CloneNo number
ARC63899

Conjugate
PE-Cy7. Ex:565nm. Em:778nm.

Background

Enables complement component C3b binding activity; heparan sulfate proteoglycan binding activity; and heparin binding activity. Contributes to cargo receptor activity. Involved in several processes, including central nervous system development; endocytosis; and positive regulation of neuron death. Acts upstream of or within several processes, including activated T cell proliferation; leukocyte migration; and microglia development. Located in external side of plasma membrane and nucleus. Is integral component of membrane. Part of integrin alphaM-beta2 complex. Is expressed in several structures, including central nervous system; heart; lymphatic vessel; thymus; and yolk sac. Human ortholog(s) of this gene implicated in lupus nephritis. Orthologous to human ITGAM (integrin subunit alpha M).

Recommended Dilutions

FC ≤0.25 µg per million cells
in 100 µl volume

Immunogen Information

Gene ID 16409	Swiss Prot P05555
-------------------------	-----------------------------

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CR3; CR3A; MAC1; Cd11b; Ly-40; Mac-1; Mac-1a; CD11b/CD18; F730045J24Rik

Contact


☎		400-999-6126
✉		cn.market@abclonal.com.cn
🌐		www.abclonal.com.cn

Product Information

Source Rabbit	Isotype IgG	Purification Affinity purification
-------------------------	-----------------------	--

Storage

Store at 2-8°C. Avoid freeze.
Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.



Flow cytometry: 1×10^6 C57BL/6 mouse bone marrow cells were surface-stained with PE/Cyanine7 Rabbit anti-Mouse CD11b mAb (A27339, 0.25 μ g, orange line) or PerCP/Cyanine5.5 Rabbit IgG isotype control (0.25 μ g, blue line). Non-fluorescently stained cells were used as blank control (red line). Viable cells were used for analysis.