

APC Rabbit anti-Mouse CD38 mAb

Catalog No.: A27365

Basic Information

Observed MW

Refer to figures

Calculated MW

34kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC59835

Conjugate

APC. Ex:650nm. Em:660nm.

Recommended Dilutions

FC ≤0.125 µg per million cells
in 100 µl volume

Background

This gene encodes a non-lineage-restricted, type II transmembrane glycoprotein that synthesizes and hydrolyzes cyclic adenosine 5'-diphosphate-ribose, an intracellular calcium ion mobilizing messenger. The release of soluble protein and the ability of membrane-bound protein to become internalized indicate both extracellular and intracellular functions for the protein. This protein has an N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites. Knockout mice deficient for this gene display altered humoral immune responses. In addition, knockout mice exhibit higher locomotor activity and defects in nurturing and social behaviors.

Immunogen Information

Gene ID

12494

Swiss Prot

P56528

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

I-19; ADPRC 1; Cd38-rs1

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1X10⁶ C57BL/6 mouse splenocytes were surface-stained with APC Rabbit IgG isotype control (A24173, 5 µl/Test, left) or APC Rabbit anti-Mouse CD38 mAb (A27365, 0.125 µg, right).