

# PerCP/Cyanine5.5 Rabbit anti-Mouse CD127/IL-7Rα mAb

Catalog No.: A27439

## **Basic Information**

#### **Observed MW**

**Calculated MW** 

51kDa

Category

Primary antibody

**Applications** 

FC

**Cross-Reactivity** 

Mouse

CloneNo number

ARC63550

Conjugate

PerCP-Cy5.5. Ex:482nm. Em:695nm.

# **Background**

Interleukin-7 is a glycoptorein involved in the regulation of lymphopoiesis. Response of cells to IL7 is dependent on the presence of the interleukin 7 receptor (IL7R); the active receptor is a alpha/gamma chain heterodimer. The gamma(c) chain, which also associates with the interleukin-2 receptor, serves primarily to activate signal transduction by the IL7R complex, while the alpha chain of IL7R determines specific signaling events through its association with cytoplasmic signaling molecules.

## **Recommended Dilutions**

FC

5  $\mu$ l per 10^6 cells in 100  $\mu$ l volume

# **Immunogen Information**

**Gene ID** 16197

**Swiss Prot** 

P16872

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

CD127; IL-7Ralpha

## **Contact**

	400-999-6126
<b>×</b>	cn.market@abclonal.com.cn
	www.abclonal.com.cn

## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification


#### Storage


Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.









Flow cytometry: 1X10^6 C57BL/6 mouse splenocytes were surface-stained with PerCP/Cyanine5.5 Rabbit IgG isotype control (A25620,5 µl/Test,blue line) or PerCP/Cyanine5.5 Rabbit anti-Mouse CD127/IL-7R $\alpha$  mAb (A27439,5  $\mu$ l/Test,orange line).Non-fluorescently stained cells were used as blank control (red line).