

PerCP/Cyanine5.5 Rabbit anti-Mouse CD127/IL-7Rα mAb

Catalog No.: A27439

Basic Information

Observed MW

Calculated MW

51kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC63550

Conjugate

PerCP-Cy5.5. Ex:482nm. Em:695nm.

Background

Interleukin-7 is a glycoptorein involved in the regulation of lymphopoiesis. Response of cells to IL7 is dependent on the presence of the interleukin 7 receptor (IL7R); the active receptor is a alpha/gamma chain heterodimer. The gamma(c) chain, which also associates with the interleukin-2 receptor, serves primarily to activate signal transduction by the IL7R complex, while the alpha chain of IL7R determines specific signaling events through its association with cytoplasmic signaling molecules.

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID 16197

Swiss Prot

P16872

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CD127; IL-7Ralpha

Contact

	400-999-6126
×	cn.market@abclonal.com.cn
	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry:1X10^6 C57BL/6 mouse splenocytes were surface-stained with ABflo® 488 anti-Mouse CD3e mAb (A26899,5 μ I/Test) and PerCP/Cyanine5.5 Rabbit IgG isotype control (A25620,5 μ I/Test,left) or PerCP/Cyanine5.5 Rabbit anti-Mouse CD127/IL-7R α mAb (A27439,5 μ I/Test,right).