Leader in Biomolecular Solutions for Life Science

www.abclonal.com

cGAS Rabbit mAb

Catalog No.: A27527 Recombinant 1 Publications

Basic Information

Observed MW
62 kDa

Calculated MW
59 kDa

Category
Primary antibody

Applications
WB,IF/ICC,IHC-P,ELISA

Cross-Reactivity
Human

CloneNo number
ARC3761

Background

Enables several functions, including 2',3'-cyclic GMP-AMP synthase activity; chromatin binding activity; and phosphatidylinositol-4,5-bisphosphate binding activity. Involved in several processes, including cellular response to exogenous dsRNA; positive regulation of intracellular signal transduction; and regulation of defense response. Located in several cellular components, including cytosol; nucleus; and site of double-strand break.

Recommended Dilutions

WB	1:1000 - 1:2000
IF/ICC	1:100 - 1:400
IHC-P	1:50 - 1:200
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
115004

Swiss Prot
Q8N884

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
MB21D1; h-cGAS; C6orf150

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20℃


Validation Data




Western blot analysis of various lysates using cGAS Rabbit mAb (A27527) at 1:1000 dilution incubated at room temperature for 1.5 hours.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Negative control (NC): Hep G2.
Exposure time: 90 s.




Confocal imaging of HT-29 cells (postive) and Hep G2 cells (negative) using cGAS Rabbit mAb (A27527, dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Immunohistochemistry analysis of paraffin-embedded Human colon tissue using cGAS Rabbit mAb (A27527) at a dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human breast tissue using cGAS Rabbit mAb (A27527) at a dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human esophagus tissue using cGAS Rabbit mAb (A27527) at a dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human lung adenocarcinoma tissue using cGAS Rabbit mAb (A27527) at a dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human lung squamous carcinoma tissue using cGAS Rabbit mAb (A27527) at a dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

Validation Data



Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using cGAS Rabbit mAb (A27527) at a dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.