APC Rabbit anti-Mouse CD155/Pvr mAb

ABclonal www.abclonal.com

Catalog No.: A27530

Basic Information

Observed MW

Calculated MW

45kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC67367

Conjugate

APC. Ex:650nm. Em:660nm.

Background

Enables dynein light chain binding activity. Acts upstream of or within cell migration and cell-cell adhesion. Located in cell surface and plasma membrane. Orthologous to human PVR (PVR cell adhesion molecule).

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID 52118

Swiss Prot

Q91WP1

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

PVS; mE4; HVED; Taa1; CD155; Tage4; necl-5; D7Ertd458e; 3830421F03Rik

Contact

2		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn


Product Information


SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at 2-8°C. Avoid freeze.


Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.







Flow cytometry: 1X10^6 NIH/3T3 cells were surface-stained with APC Rabbit IgG isotype control (A24173,5 μ I/Test,Ieft) or APC Rabbit anti-Mouse CD155/Pvr mAb (A27530,5 μ I/Test,right)



Flow cytometry: 1X10^6 C57BL/6 mouse splenocytes were surface-stained with APC Rabbit anti-Mouse CD155/Pvr mAb (A27530,5 µl/Test,orange line) or APC Rabbit IgG isotype control (A24173,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).