ABclonal www.abclonal.com

ABflo® 488 Rabbit anti-Human CD45RB mAb

Catalog No.: A27566

Basic Information

Observed MW

Calculated MW

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC65145

Conjugate

ABflo® 488. Ex:491nm. Em:516nm.

Background

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitosis, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported.

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID 5788 **Swiss Prot**

P08575-9

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 26-463 of human CD45RB (XP_006711537.1).

Synonyms

LCA; LY5; B220; CD45; L-CA; T200; CD45R; GP180; IMD105

Contact

2		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data









Flow cytometry: 1X10^6 MOLT-4 cells were

surface-stained with ABflo® 488 Rabbit IgG


ABflo® 488 Rabbit anti-Human CD45RB mAb

isotype control (A22069,5 µl/Test,left) or






Flow cytometry: 1X10^6 HAP1 cells (negative control,left) and MOLT-4 cells (right) were surface-stained with ABflo® 488 Rabbit anti-Human CD45RB mAb (A27566,5 µl/Test,orange line) or ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1X10^6 HAP1 cells (negative control,left) and Human PBMC (right) were surface-stained with ABflo® 488 Rabbit anti-Human CD45RB mAb (A27566,5 µl/Test,orange line) or ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line). Cells in the lymphocyte gate were used for analysis.



Flow cytometry: 1X10^6 Human PBMC were surface-stained with ABflo® 488 Rabbit IgG isotype control (A22069,5 μ I/Test,left) or ABflo® 488 Rabbit anti-Human CD45RB mAb (A27566,5 μ I/Test,right). Cells in the lymphocyte gate were used for analysis.