# APC Rat anti-Mouse CD45R/B220 mAb

www.abclonal.com

ABclonal

Catalog No.: A27570

## **Basic Information**

#### **Observed MW**

Calculated MW

130kDa/140kDa/144kDa

Category

Primary antibody

**Applications** 

FC

**Cross-Reactivity** 

Mouse

CloneNo number

ATC10005

Conjugate

APC. Ex:650nm. Em:660nm.

# **Background**

Enables several functions, including heparan sulfate proteoglycan binding activity; heparin binding activity; and protein tyrosine phosphatase activity. Involved in several processes, including lymphocyte differentiation; positive regulation of macromolecule metabolic process; and regulation of signal transduction. Acts upstream of or within several processes, including lymphocyte differentiation; positive regulation of lymphocyte activation; and regulation of protein phosphorylation. Located in external side of plasma membrane; focal adhesion; and membrane raft. Is expressed in several structures, including 3rd branchial arch; alimentary system; cardiovascular system; hemolymphoid system; and placenta. Used to study systemic lupus erythematosus. Human ortholog(s) of this gene implicated in hepatitis C; multiple sclerosis; severe combined immunodeficiency; and severe combined immunodeficiency, autosomal recessive, T cell-negative, B cell-positive, Nk cell-positive. Orthologous to human PTPRC (protein tyrosine phosphatase receptor type C).

### **Recommended Dilutions**

FC

5  $\mu$ l per 10^6 cells in 100  $\mu$ l volume

### **Immunogen Information**

**Gene ID** 19264

**Swiss Prot** 

P06800

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

loc; B220; Cd45; L-CA; Ly-5; T200; CD45R; Lyt-4

### **Contact**

2		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

### **Product Information**


SourceIsotypePurificationRatIgGAffinity purification

### **Storage**

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

# **Validation Data**



Flow cytometry:1X10^6 C57BL/6 mouse splenocytes were surface-stained with ABflo® 450 Rabbit anti-Mouse CD19 (A26192,5 µl/Test) and APC Rat IgG2a isotype control (5 µl/Test,left) or APC Rat anti-Mouse CD45R/B220 mAb (A27570,5 µl/Test,right).