APC anti-Mouse TCR γ/δ mAb

Catalog No.: A27574



Basic Information

Observed MW

Calculated MW

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ACC0007

Conjugate

APC. Ex:650nm. Em:660nm.

Background

T cell receptors recognize foreign antigens which have been processed as small peptides and bound to major histocompatibility complex (MHC) molecules at the surface of antigen presenting cells (APC). Each T cell receptor is a dimer consisting of one alpha and one beta chain or one delta and one gamma chain. In a single cell, the T cell receptor loci are rearranged and expressed in the order delta, gamma, beta, and alpha. If both delta and gamma rearrangements produce functional chains, the cell expresses delta and gamma. If not, the cell proceeds to rearrange the beta and alpha loci. This region represents the germline organization of the T cell receptor alpha and delta loci. Both the alpha and delta loci include V (variable), J (joining), and C (constant) segments and the delta locus also includes diversity (D) segments. The delta locus is situated within the alpha locus, between the alpha variable and joining genes. During T cell development, the delta chain is synthesized by a recombination event at the DNA level joining a D segment with a I segment; a V segment is then joined to the D-J gene. The alpha chain is synthesized by recombination joining a single V segment with a J segment. For both chains, the C segment is later joined by splicing at the RNA level. Recombination of many different V segments with several | segments provides a wide range of antigen recognition. Additional diversity is attained by junctional diversity, resulting from the random additional of nucleotides by terminal deoxynucleotidyltransferase. Ten variable segments can be used in either alpha or delta chains and are described by TRAV/DV symbols. Several V and J segments of the alpha locus are known to be incapable of encoding a protein and are considered pseudogenes.

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID

Swiss Prot

110066/110067

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

Tcrdelta; TCRGV1S1; TCRGV2S1; TCRGV3S1; TCRGV5S3

Contact

a	400-999-6126
\sim	cn.market@abclonal.com.cn
⊙	www.abclonal.com.cn

Product Information


SourceIsotypePurificationArmenian HamsterArmenian Hamster IgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry:1X10^6 C57BL/6 mouse splenocytes were surface-stained with ABflo® 488 Rat anti-Mouse CD3 mAb (A27161,5 μ l/Test) and APC Armenian Hamster IgG isotype control (5 μ l/Test,left) or APC anti-Mouse TCR γ / δ mAb (A27574,5 μ l/Test,right). Cells in the lymphocyte gate were used for analysis.