Leader in Biomolecular Solutions for Life Science

www.abclonal.com

ABflo® 680 Rabbit IgG isotype control

Catalog No.: A27697

Basic Information

Observed MW

Calculated MW

Category
Primary antibody

Applications
FC

Cross-Reactivity

CloneNo number
ARC5105-10

Conjugate
ABflo® 680. Ex:679nm. Em:700nm.

Background

The isotype of a primary antibody and the application it is being used in can result in background staining. Primary antibody background noise can be caused by binding to Fc receptors on target cells; by non-specific interactions with cellular proteins, carbohydrates, and lipids; or by cell autofluorescence. Isotype control antibodies can act as negative controls to help differentiate non-specific background signal from specific antibody signal because they have no relevant specificity to a target antigen.An isotype control antibody should have the same immunoglobulin type and be used at the same concentration as the test antibody.

Recommended Dilutions

FC	≤0.25 μg per million cells in 100 μl volume

Immunogen Information

Gene ID

Swiss Prot

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20℃

Validation Data



Flow cytometry: 1X10^6 293F cells (negative control,left) and Human PBMC (right) were surface-stained with ABflo® 680 Rabbit anti-Human CD8a mAb (5 µl/Test,orange line) or ABflo® 680 Rabbit IgG isotype control (A27697,0.25µg,blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1X10^6 Human PBMC were surface-stained with ABflo® 680 Rabbit IgG isotype control (A27697,0.25µg,left) or ABflo® 680 Rabbit anti-Human CD8a mAb (5 µl/Test,right).