

# ABflo® 450 Rat anti-Mouse MHC Class II (I-A/I-E) mAb

**Catalog No.: A27805**

## Basic Information

**Observed MW****Calculated MW**

26kDa/30kDa/32kDa

**Category**

Primary antibody

**Applications**

FC

**Cross-Reactivity**

Mouse

**CloneNo number**

ATC50002

**Conjugate**

ABflo® 450. Ex:406nm. Em:445nm.

## Background

Predicted to enable several functions, including MHC class II receptor activity; peptide antigen binding activity; and signaling receptor binding activity. Involved in response to interferon-gamma. Acts upstream of or within antigen processing and presentation of exogenous peptide antigen via MHC class II. Predicted to be located in external side of plasma membrane and immunological synapse. Predicted to be part of MHC class II protein complex. Predicted to be active in late endosome membrane and lysosomal membrane. Is expressed in lung. Human ortholog(s) of this gene implicated in several diseases, including asthma (multiple); autoimmune disease (multiple); bacterial infectious disease (multiple); eye disease (multiple); and sarcoidosis (multiple). Orthologous to human HLA-DRB1 (major histocompatibility complex, class II, DR beta 1) and HLA-DRB5 (major histocompatibility complex, class II, DR beta 5).

## Recommended Dilutions

**FC** ≤0.5 µg per million cells  
in 100 µl volume

## Immunogen Information

**Gene ID**

14969/14961/381091/14960

**Swiss Prot**

P18468/P06342/Q3U060/P14435

**Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

**Synonyms**

Eb; Ia4; H2Eb; Ia-4; H-2Eb;IAb; Ia2; Ia-2; Abeta; H-2Ab; H2-Ab; Rmcs1; I-Abeta; Ia5; Ia-5; H-2Eb2; Iebeta2; A130038H09Rik; Ia1; H2Aa; Ia-1; H-2Aa; Aalpha; IAalpha; I-Aalpha

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

**Source**

Rat

**Isotype**

IgG

**Purification**


Affinity purification

**Storage**


Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

# Validation Data



Flow cytometry:  $1 \times 10^6$  Neuro-2a cells (negative control, left) and C57BL/6 mouse splenocytes (right) were surface-stained with ABflo® 450 Rat anti-Mouse MHC Class II (I-A/I-E) mAb (A27805, 0.5  $\mu$ g, orange line) or ABflo® 450 Rat IgG isotype control (0.5  $\mu$ g, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry:  $1 \times 10^6$  C57BL/6 mouse splenocytes were surface-stained with ABflo® 450 Rat IgG isotype control (0.5  $\mu$ g, left) or ABflo® 450 Rat anti-Mouse MHC Class II (I-A/I-E) mAb (A27805, 0.5  $\mu$ g, right).