




## Validation Data



Flow cytometry:  $1 \times 10^6$  Human PBMC (untreated, left) and Human PBMC (treated with 50 ng/ml PMA and 1  $\mu$ g/ml Ionomycin for 6 hours and 2  $\mu$ M Monensin for last 5 hours, right) were surface-stained with ABflo® 450 Rabbit anti-Human/Monkey CD3 mAb (A27177, 5  $\mu$ l/Test), and then intracellularly-stained with APC/Cyanine7 Rabbit anti-Human IFN-gamma mAb (A27830, 5  $\mu$ l/Test). Cells in the lymphocyte gate were used for analysis.

Flow cytometry:  $1 \times 10^6$  Cynomolgus PBMC (treated with 50 ng/mL PMA and 1  $\mu$ g/mL Ionomycin for 4-6 hours in the presence of monensin) were surface-stained with ABflo® 450 Rabbit anti-Human/Monkey CD3 mAb (A27177, 5  $\mu$ l/Test) and then intracellularly-stained with APC/Cyanine7 Rabbit IgG isotype control (5  $\mu$ l/Test, left) or APC/Cyanine7 Rabbit anti-Human/Monkey IFN-gamma mAb (A27830, 5  $\mu$ l/Test, right). Cells in the lymphocyte gate were used for analysis.