

# Pit1/POU1F1 Rabbit mAb

Catalog No.: A27875 **Recombinant**

## Basic Information

### Observed MW

33kDa

### Calculated MW

33kDa

### Category

Primary antibody

### Applications

WB,IP,IF/ICC,IHC-P,ELISA

### Cross-Reactivity

Mouse, Rat

### CloneNo number

ARC73912

## Background

This gene encodes a member of the POU family of transcription factors that regulate mammalian development. The protein regulates expression of several genes involved in pituitary development and hormone expression. Mutations in this genes result in combined pituitary hormone deficiency. Multiple transcript variants encoding different isoforms have been found for this gene.

## Recommended Dilutions

**WB** 1:4000 - 1:16000

**IP** 0.5µg-4µg antibody for  
200µg-400µg extracts of  
whole cells

**IF/ICC** 1:200 - 1:800

**IHC-P** 1:1000 - 1:4000

**ELISA** Recommended starting  
concentration is 1 µg/mL.  
Please optimize the  
concentration based on  
your specific assay  
requirements.

## Immunogen Information

### Gene ID

5449

### Swiss Prot

P28069

### Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

### Synonyms

PIT1; CPHD1; GHF-1; Pit-1; POU1F1a

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


## Contact

☎ | 400-999-6126


✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)


Validation Data



Immunoprecipitation of Pit1/POU1F1 from 300 µg extracts of Mouse pituitary tissue was performed using 2 µg of Pit1/POU1F1 Rabbit mAb (A27875). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using Pit1/POU1F1 Rabbit mAb (A27875) at a dilution of 1:9000.




Western blot analysis of various lysates using Pit1/POU1F1 Rabbit mAb (A27875) at 1:8000 dilution incubated overnight at 4°C.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25 µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Negative control (NC): Mouse brain  
Exposure time: 10s.




Western blot analysis of various lysates using Pit1/POU1F1 Rabbit mAb (A27875) at 1:8000 dilution incubated overnight at 4°C.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25 µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Negative control (NC): Rat liver  
Exposure time: 60s.


## Validation Data




Confocal imaging of paraffin-embedded Rat pituitary tissue using Pit1/POU1F1 Rabbit mAb (A27875, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Confocal imaging of paraffin-embedded Mouse pituitary tissue using Pit1/POU1F1 Rabbit mAb (A27875, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Immunohistochemistry analysis of paraffin-embedded Rat pituitary gland tissue using Pit1/POU1F1 Rabbit mAb (A27875) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse pituitary gland tissue using Pit1/POU1F1 Rabbit mAb (A27875) at a dilution of 1:4000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.