# PE/Cyanine7 Rabbit anti-Human CD27 mAb

ABclonal www.abclonal.com

Catalog No.: A27909

## **Basic Information**

#### **Observed MW**

**Calculated MW** 

29kDa

Category

Primary antibody

**Applications** 

FC

**Cross-Reactivity** 

Human

CloneNo number

ARC54542

Conjugate

PE-Cy7. Ex:565nm. Em:778nm.

## **Background**

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. This receptor transduces signals that lead to the activation of NF-kappaB and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA), a proapoptotic protein, can bind to this receptor and is thought to play an important role in the apoptosis induced by this receptor.

## **Recommended Dilutions**

FC

5  $\mu$ l per 10^6 cells in 100  $\mu$ l volume

## Immunogen Information

Gene ID 939 Swiss Prot

P26842

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

T14; S152; Tp55; TNFRSF7; S152. LPFS2

## **Contact**

2		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### **Storage**


Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

## **Validation Data**







Flow cytometry: 1X10^6 HEL cells (negative control,left) and Daudi cells (right) were surface-stained with PE/Cyanine7 Rabbit anti-Human CD27 mAb (A27909,5 µl/Test,orange line) or PE/Cyanine7 Rabbit IgG isotype control (5 µl/Test,blue line). Nonfluorescently stained cells were used as blank control (red line).

Flow cytometry: 1X10^6 Human PBMC were surface-stained with PE/Cyanine7 Rabbit anti-Human CD27 mAb (A27909,5  $\mu$ I/Test,orange line) or PE/Cyanine7 Rabbit IgG isotype control (5  $\mu$ I/Test,blue line). Nonfluorescently stained cells were used as blank control (red line).