






## Validation Data



Flow cytometry:  $1 \times 10^6$  Cynomolgus PBMC were surface-stained with ABflo® 450 Rabbit anti-Human/Monkey CD19 mAb (A27286, 5  $\mu\text{l}/\text{Test}$ ) and APC/Cyanine7 Rabbit IgG isotype control (5  $\mu\text{l}/\text{Test}$ , left) or APC/Cyanine7 Rabbit anti-NHP IgD mAb (A27949, 5  $\mu\text{l}/\text{Test}$ , right). Cells in the lymphocyte gate were used for analysis.



Flow cytometry:  $1 \times 10^6$  Rhesus PBMC were surface-stained with ABflo® 450 Mouse anti-Human CD11b mAb (A27475, 5  $\mu\text{l}/\text{Test}$ ), ABflo® 488 Rabbit anti-Human/Monkey CD19 mAb (A23008, 5  $\mu\text{l}/\text{Test}$ ) and APC/Cyanine7 Rabbit IgG isotype control (5  $\mu\text{l}/\text{Test}$ , left) or APC/Cyanine7 Rabbit anti-NHP IgD mAb (A27949, 5  $\mu\text{l}/\text{Test}$ , right). Cells in the CD11b- gate were used for analysis.



Flow cytometry:  $1 \times 10^6$  Human peripheral blood mononuclear cells (negative control, left) and Cynomolgus peripheral blood mononuclear cells (right) were surface-stained with APC/Cyanine7 Rabbit anti-NHP IgD mAb (A27949, 5  $\mu\text{l}/\text{Test}$ , right). Cells in the lymphocyte gate were used for analysis.