

APC/Cyanine7 Rabbit anti-Human/Monkey HLA-DR mAb

Catalog No.: A27950

Basic Information

Observed MW**Calculated MW**

29kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human, Cynomolgus

CloneNo number

ARC5141-01

Conjugate

APC-Cy7. Ex:651nm. Em:779nm.

Background

HLA-DRA is one of the HLA class II alpha chain paralogues. This class II molecule is a heterodimer consisting of an alpha and a beta chain, both anchored in the membrane. This molecule is expressed on the surface of various antigen presenting cells such as B lymphocytes, dendritic cells, and monocytes/macrophages, and plays a central role in the immune system and response by presenting peptides derived from extracellular proteins, in particular, pathogen-derived peptides to T cells. The alpha chain is approximately 33-35 kDa and its gene contains 5 exons. Exon 1 encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, and exon 4 encodes the transmembrane domain and the cytoplasmic tail. DRA does not have polymorphisms in the peptide binding part and acts as the sole alpha chain for DRB1, DRB3, DRB4 and DRB5.

Recommended Dilutions

FC 5 µl per 10⁶ cells in
100 µl volume

Immunogen Information

Gene ID

3122

Swiss Prot

P01903

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

HLA-DRA1

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.09% Sodium azide, 0.2% BSA, pH7.3.

Validation Data



Flow cytometry: 1X10⁶ Human PBMC were surface-stained with ABflo® 488 Rabbit anti-Human/Monkey CD14 mAb (A25071, 5 µl/Test) and APC/Cyanine7 Rabbit IgG isotype control (5 µl/Test, left) or APC/Cyanine7 Rabbit anti-Human/Monkey HLA-DR mAb (A27950, 5 µl/Test, right). Cells in the lymphocyte and monocyte gate were used for analysis.