

[KO Validated] ATF3 Rabbit mAb

Catalog No.: A28345 **KO** **Validated** **Recombinant**

Basic Information

Observed MW

23 kDa/24 kDa

Calculated MW

12-21 kDa

Category

Primary antibody

Applications

WB,IP,ChIP,ELISA

Cross-Reactivity

Human, Mouse

CloneNo number

ARC77618

Background

This gene encodes a member of the mammalian activation transcription factor/cAMP responsive element-binding (CREB) protein family of transcription factors. This gene is induced by a variety of signals, including many of those encountered by cancer cells, and is involved in the complex process of cellular stress response. Multiple transcript variants encoding different isoforms have been found for this gene. It is possible that alternative splicing of this gene may be physiologically important in the regulation of target genes.

Recommended Dilutions

WB 1:2000 - 1:5000**IP** 0.5µg-4µg antibody for
200µg-400µg extracts of
whole cells**ChIP** 5µg antibody for 10µg-15µg
of Chromatin**ELISA** Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on your
specific assay requirements.
For high-ratio antibody
dilutions (≥1:10000) a
sequential dilution method
is strongly recommended to
ensure measurement
accuracy.

Immunogen Information

Gene ID

467

Swiss Prot

P18847

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

ATF3

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data



Immunoprecipitation of ATF3 from 300 µg extracts of 293T cells was performed using 2 µg of [KO Validated] ATF3 Rabbit mAb (A28345). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1x reducing Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using [KO Validated] ATF3 Rabbit mAb (A28345) at a dilution of 1:5000.




Western blot analysis of lysates from wild type (WT) and ATF3 knockout (KO) 293T cells using [KO Validated] ATF3 Rabbit mAb (A28345) at 1:5000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90 s.



Western blot analysis of various lysates using [KO Validated] ATF3 Rabbit mAb (A28345) at 1:5000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90 s.

Validation Data



Chromatin immunoprecipitation was performed with 15 µg of cross-linked chromatin from 293T cells, using 5 µg of [KO Validated] ATF3 Rabbit mAb (A28345) and Rabbit IgG isotype control (AC042). The enrichment of immunoprecipitated DNA at different genomic loci was examined by quantitative PCR. The histogram compares the ratio of the immunoprecipitated DNA to the input at given loci.